















Dressing Name	Product Images	Description / Type of Dressing	Use in Radiotherapy	How to Apply	Size Available
<p><b>PolyMem®</b> <b>PolyMem® MAX</b></p>		<p><b>PolyMem:</b> Absorbent, hydrophilic polyurethane foam containing a cleanser and moisturiser with semipermeable film backing.</p> <p><b>PolyMem MAX:</b> As above but 60% more absorbent.</p> <p><b>Polymem Roll:</b> As Polymem but on a roll to cut to size.</p>	<ul style="list-style-type: none"> <li>- Brisk erythema, dry and moist desquamation (RTOG2, 2.5 and 3).</li> <li>- Non - low exuding areas (Polymem, Polymem Roll).</li> <li>- Moderate - high exuding areas (Polymem Max, Polymem Roll).</li> <li>- Malignant, fungating wounds.</li> </ul>	<p>Apply directly to wound with an overlap margin of between 0.6 - 5cm.</p> <p>Apply with printed side outwards.</p> <p>Can be cut to shape and size.</p> <p>Dressing change dependent upon amount of exudate.</p> <p>Wound / radiotherapy field does not require cleansing at dressing changes unless area infected, contaminated or excess amounts of exudate present.</p> <p>Will need additional fixation i.e. Mepitac, Tubifast, Fixation pants, Tracheostomy tube velcro straps (neck collar).</p> <p>Can be moistened (sprinkled) with water prior to applying to dry desquamation to soothe and moisturise area.</p>	<p><b>PolyMem:</b> 8cm x 8cm 10cm x 10cm 13cm x 13cm 17cm x 19cm 10cm x 61cm roll 20cm x 60cm roll</p> <p><b>PolyMem MAX:</b> 11cm x 11cm 20cm x 20cm</p>
<p><b>PolyMem® Tube</b></p>		<p><b>PolyMem Tube:</b> Absorbent, hydrophilic polyurethane foam containing a cleanser and moisturiser with semipermeable film backing. Pre-cut for fitting around tube sites</p>	<ul style="list-style-type: none"> <li>- Around tube sites (eg. PEG or Tracheostomy sites) that are in the field of radiotherapy for management of skin reactions.</li> <li>- Around tube sites in locations outside of the field of radiotherapy for management of skin complications around the tube site e.g. Hypergranulation, pressure and friction damage.</li> </ul>	<p>Dressings are pre-cut to fit around the tube sites.</p> <p>For Tracheostomy sites that are sewn in place, the dressing may need to be cut to a narrower width to insert under the stitched-in cuff.</p> <p>The tube holders and straps will usually hold the dressing in place, however additional fixation may be required as necessary.</p>	<p><b>PolyMem Tube:</b> 7cm x 7cm 9cm x 9cm</p>
<p><b>PolyMem® WIC</b></p>		<p><b>PolyMem WIC:</b> Perforated, absorbent, hydrophilic polyurethane foam containing cleanser and moisturiser without film backing.</p>	<ul style="list-style-type: none"> <li>- Brisk erythema, dry and moist desquamation (RTOG 2, 2.5 and 3).</li> <li>- Cavity wounds.</li> <li>- Between skin folds in treatment field i.e. buttocks, groins, inframammary fold, axilla.</li> </ul>	<p>Can be cut to shape and size.</p> <p>Cleanses and moisturises all surfaces in contact with the dressing due to absence of film backing.</p> <p>Place as needed between skin folds to cleanse, soothe and moisturise promoting patient comfort.</p> <p>Secondary dressing / absorbent pad required / fixation pants required. Allows exudate to transfer through to secondary dressing / absorbent pad.</p> <p>When used in cavities ensure filler is trimmed to 1/3rd smaller than open cavity and place in gently (do not pack tightly / overfill).</p> <p>Cover area with a secondary dressing.</p>	<p><b>PolyMem WIC:</b> 8cm x 8cm</p>

Dressing Name	Product Images	Description / Type of Dressing	Use in Radiotherapy	How to Apply	Size Available
<ul style="list-style-type: none"> <li>● Mepilex®</li> <li>● Mepilex® Lite</li> </ul>		<ol style="list-style-type: none"> <li>1. Flexible, absorbent, polyurethane foam dressing with a unique Safetac® soft silicone wound contact layer and a vapour permeable, waterproof outer film.</li> <li>2. 'Lite' is a thinner, more flexible version of Mepilex.</li> </ol>	<ul style="list-style-type: none"> <li>- Dry and moist desquamation (RTOG 2, 2.5 and 3 reactions).</li> <li>- Non - low exuding wounds (Lite).</li> <li>- Low to moderate exuding wounds (Mepilex).</li> <li>- 'Lite' useful for protection of fragile skin and minimising friction in skin folds.</li> </ul>	Dressing to overlap area being covered by 2 – 3 cms. Can be cut to shape and size. Secondary dressing not required but if additional fixation needed consider Mepitac / Tubifast / Fixation pants (pelvic radiotherapy fields). Dressing change dependent upon amount of exudate. Change when strikethrough within 1 – 2 cms of dressing edges.	<p><b>Mepilex:</b> 10cm x 11cm, 11cm x 20cm 15cm x 16cm, 20cm x 21cm 20cm x 50cm</p> <p><b>Mepilex Lite:</b> 6cm x 8.5cm, 10cm x 10cm, 15cm x 15cm 20cm x 50cm</p>
<ul style="list-style-type: none"> <li>● Mepilex® Border</li> <li>● Mepilex® Border Lite</li> </ul>		<ol style="list-style-type: none"> <li>1. Flexible, absorbent polyurethane foam island dressing with a unique Safetac® soft silicone wound contact layer and border. The outer film is vapour permeable and waterproof.</li> <li>2. 'Lite' version is less absorbent.</li> </ol>	<ul style="list-style-type: none"> <li>- Moderate - high exuding areas of moist desquamation (RTOG 2.5 and 3).</li> <li>- Basal Cell Carcinoma (BCC) skin lesions.</li> <li>- Malignant, fungating wounds.</li> </ul>	Apply directly to wound ensuring border is onto intact, non-friable skin. Dressing change dependent upon amount of exudate. Change when strikethrough within 1 – 2 cms of of foam island edges.	<p><b>Mepilex Border:</b> 7cm x 7.5cm, 10cm x 12.5cm 15cm x 17.5cm, 17cm x 20cm 10cm x 20cm, 10cm x 30cm</p> <p><b>Mepilex Border Lite:</b> 4cm x 5cm, 5cm x 12.5cm, 7.5cm x 7.5cm, 10cm x 10cm, 15cm x 15cm</p>
<ul style="list-style-type: none"> <li>● Mepitel®</li> <li>● Mepitel® One</li> </ul>		<ol style="list-style-type: none"> <li>1. Porous, transparent, flexible wound contact layer consisting of a polyimide net coated on both sides with a unique Safetac® soft silicone adhesive technology.</li> <li>2. Transparent, flexible wound contact layer consisting of a perforated polyurethane film coated on one side with unique Safetac® soft silicone.</li> </ol>	<ul style="list-style-type: none"> <li>- Dry and moist desquamation.</li> <li>- Malignant, fungating wounds.</li> <li>- To protect fragile skin.</li> <li>- BCC skin lesions.</li> <li>- Cavity wounds i.e. orbital / nasal areas.</li> </ul>	Place gently and loosely into cavity. Can be used with a hydrogel. Can be cut to shape and size. Requires appropriate secondary dressing.	<p><b>Mepitel:</b> 5cm x 7cm, 8cm x 10cm, 12cm x 15cm, 20cm x 32cm</p> <p><b>Mepitel One:</b> 6cm x 7cm, 9cm x 10cm, 13cm x 15cm, 24cm x 27.5cm</p>
<ul style="list-style-type: none"> <li>● Mepilex® Transfer</li> </ul>		Exudate transfer dressing consisting of a thin, very flexible polyurethane foam with unique Safetac® soft silicone.	<ul style="list-style-type: none"> <li>- Moderate - high exuding areas of moist desquamation (RTOG 2.5 and 3).</li> <li>- Difficult to dress areas particularly groins, genital and anal areas in patients receiving pelvic radiotherapy.</li> </ul>	Dressing needs to overlap onto dry surrounding skin by approx 2 – 3 cms. Secondary absorbent dressing required. Can be cut to shape and size.	<p><b>Mepilex Transfer:</b> 7.5cm x 8.5cm, 10cm x 12cm 15cm x 20cm, 20cm x 50cm</p>
<ul style="list-style-type: none"> <li>● Mepitac®</li> </ul>		Conformable tape with unique Safetac® soft silicone.	Fixation of dressings within radiotherapy treatment field.	Apply to edges of dressing that needs to be secured / requires extra fixation, positioning tape onto intact skin.	<p><b>Mepitac:</b> 2cm x 300cm, 4cm x 150cm</p>

<i>Dressing Name</i>	<i>Product Images</i>	<i>Description / Type of Dressing</i>	<i>Use in Radiotherapy</i>	<i>How to Apply</i>	<i>Size Available</i>
<b>ALLEVYN® GENTLE</b>		Absorbent, hydrocellular polyurethane foam dressing consisting of a perforated soft gel adhesive wound contact layer and a highly permeable waterproof outer layer.	<ul style="list-style-type: none"> <li>- Moderate - high exuding areas of moist desquamation (RTOG 2.5 and 3).</li> <li>- Malignant, fungating wounds.</li> </ul>	Apply directly to skin reaction - white side to skin. Requires additional fixation to hold in place i.e. Mepitac / Tubifast / Fixation pants (pelvic radiotherapy fields). Can be cut to shape and size. Dressing change dependent upon amount of exudate.	<b>Allevyn Gentle:</b> 5cm x 5cm 10cm x 10cm 15cm x 15cm 10cm x 20cm 20cm x 20cm
<b>ALLEVYN GENTLE BORDER</b>		Adhesive bordered, hydrocellular polyurethane foam pad dressing consisting of a perforated soft gel wound contact layer and highly permeable waterproof outer layer.	<ul style="list-style-type: none"> <li>- Moderate - high exuding wounds (RTOG 2.5 and 3).</li> <li>- Malignant, fungating wounds.</li> <li>- BCC skin lesions.</li> </ul>	Apply directly to wound ensuring adhesive border is onto intact, non-friable skin. Dressing change dependent upon amount of exudate.	<b>Allevyn Gentle Border:</b> 7.5cm x 7.5cm 10cm x 10cm 12.5cm x 12.5cm 17.5cm x 17.5cm
<b>ALLEVYN LITE</b>		Thin, absorbent, hydrocellular polyurethane foam dressing consisting of a perforated non-adherent wound contact layer and a waterproof outer film	<ul style="list-style-type: none"> <li>- Low - moderate exuding wounds</li> <li>- Dry and patchy moist desquamation (RTOG 2 and 2.5)</li> <li>- Between skin folds to prevent skin on skin friction</li> <li>- Neck collar</li> </ul>	Apply directly to skin reaction - white side to skin. Requires additional fixation with silicone tape / tracheostomy tube velcro straps(collar) / fixation pants. Can be cut to shape and size. Dressing change dependent upon amount of exudate.	<b>Allevyn Lite:</b> 5cm x 5cm 10cm x 10cm 10cm x 20cm 15cm x 20cm
<b>ALLEVYN</b>		Absorbent hydrocellular polyurethane foam dressing consisting of a perforated non-adherent wound contact layer and a waterproof outer film	<ul style="list-style-type: none"> <li>- Moderate - high exuding wounds</li> <li>- Patchy and confluent moist desquamation (RTOG 2.5 and 3).</li> <li>- Neck Collar</li> </ul>	Apply directly to skin reaction - white side to skin. Requires additional fixation with silicone tape / tracheostomy tube velcro straps(collar) / fixation pants. Can be cut to shape and size. Dressing change dependent upon amount of exudate.	<b>Allevyn Lite:</b> 5cm x 5cm 10cm x 10cm 10cm x 20cm 20cm x 20cm
<b>ALLEVYN GENTLE BORDER LITE</b>		Adhesive bordered hydrocellular dressing with a slimmer foam core consisting of a perforated soft gel wound contact layer and highly permeable waterproof outer layer.	<ul style="list-style-type: none"> <li>- Low - moderate exuding wounds</li> <li>- Patchy moist desquamation (RTOG 2.5)</li> <li>- Malignant, fungating wounds</li> <li>- BCC skin lesions</li> </ul>	Apply directly to wound ensuring adhesive border is onto intact, non-friable skin. Dressing change dependent upon amount of exudate.	<b>Allevyn Gentle Border Lite:</b> 5cm x 5cm 7.5cm x 7.5cm 10cm x 10cm 5.5cm x 12.5cm 8cm x 15cm 15cm x 15cm
<b>ALLEVYN HEEL</b>		'Cup shaped' absorbent hydrocellular polyurethane foam dressing consisting of a perforated non-adherent wound contact layer and a waterproof outer film.	Moderate - high exuding wounds Patchy / confluent moist desquamation (RTOG 2.5 and 3) Difficult to dress areas i.e. scrotum / penis	Apply directly to skin reaction - white side to skin. Requires additional fixation i.e. fixation pants. Dressing change dependent upon amount of exudate.	<b>Allevyn Heel:</b> 10.5cm x 13.5cm